



International Journal of Applied Sciences and Biotechnology

A Rapid Publishing Journal

ISSN 2091-2609

Indexing and Abstracting

CrossRef, Google Scholar, Global Impact Factor, Genamics, Index Copernicus, Directory of Open Access Journals, WorldCat, Electronic Journals Library (EZB), Universitätsbibliothek Leipzig, Hamburg University, UTS (University of Technology, Sydney): Library, International Society of Universal Research in Sciences (EyeSource), Journal Seeker, WZB, Socolar, BioRes, Indian Science, Jadoun Science, Jour-Informatics, Journal Directory, JournalTOCs, Academic Journals Database, Journal Quality Evaluation Report, PDOAJ, Science Central, Journal Impact Factor, NewJour, Open Science Directory, Directory of Research Journals Indexing, Open Access Library, International Impact Factor Services, SciSeek, Cabell's Directories, Scientific Indexing Services, CiteFactor, UniSA Library, InfoBase Index, Infomine, Getinfo, Open Academic Journals Index, HINARI, etc.

CODEN (Chemical Abstract Services, USA): IJASKD

Vol-3(4) December, 2015

Available online at:

<http://www.ijasbt.org>

&

<http://www.nepjol.info/index.php/IJASBT/index>



Impact factor*: 1.422

Scientific Journal Impact factor#: 3.419

Index Copernicus Value: 6.02

IBI Factor 2015**: 4.19

*Impact factor is issued by Universal Impact Factor. Kindly note that this is not the IF of Journal Citation Report (JCR).

#Impact factor is issued by SJIF INNO SPACE; **Impact factor is issued by INFOBASE INDEX.

For any type of query and/or feedback don't hesitate to email us at: editor.ijasbt@gmail.com



Short Communication

CECOCOLIC INTUSSUSCEPTION IN AN ASIAN ELEPHANT (*ELEPHAS MAXIMUS*) IN SRI LANKA ASSOCIATED WITH CHRONIC HEPATO-INTESTINAL SCHISTOSOMIASIS: FIRST CASE REPORT

A.C. Karawita*, R.C. Rajapakse, P.G.I.D. Amarasiri, R.M.T.M. Ramanayake, W.R. Jayaweera, Trent K. Bollinger and G.S.P.De.S. Gunawardena

Department of Veterinary Pathobiology, Faculty of Veterinary Medicine & Animal Science, University of Peradeniya, Peradeniya, 20400, Sri Lanka

*Corresponding author's email: ackarawita@yahoo.com

Abstract

A 25 year old female captive Asian elephant weighing approximately 3000kgs died and was subjected to a complete necropsy within 5 hours *post mortem*. Grossly, the elephant had sub-cutaneous edema, cecocolic intussusception and ascites. Histopathological examination revealed multifocal, granulomatous, portal phlebitis in the liver, multifocal, granulomatous, peri-portal hepatitis with marked portal and perisinusoidal fibrosis, and multifocal granulomatous colitis with intralesional *Schistosoma* eggs. To our knowledge, this is the first report of cecocolic intussusception in an Asian elephant associated with *Schistosoma* infection.

Keywords: *Elephas maximus*; Asian elephant; Intussusception; Schistosomiasis; Sri Lanka; *Bivitellobilharzia nairi*

Case Report

A complete necropsy was carried out within 5 hours *post mortem* on a 25 year old female Asian elephant weighing approximately 3000kgs living in the Elephant Orphanage in Pinnawala, Sri Lanka (7.3006° N, 80.3883° E). The animal had died despite treatment for chronic indigestion and progressive emaciation of approximately one month.

Sections from all the major organs and representative samples from conspicuous lesions were fixed in 10% neutral buffered formalin, dehydrated in a graded series of ethanol to xylene, embedded in paraffin and sectioned at 4 micrometers. The sections were stained with hematoxylin and eosin (HE) and Masson's Trichrome Stain.

Macroscopically, there were no visible subcutaneous or visceral fat reserves. The orbital fossae were depressed and the wings of both ilea were prominent, all of which are indicative of poor nutritional condition (Fig. 1a). The elephant had died in lateral recumbancy and there was marked diffuse subcutaneous edema of the ventral body wall. The peritoneal cavity contained over five liters of yellow turbid fluid (Fig. 1b). The liver was markedly congested and diffusely enlarged; there were multifocal to coalescing white foci on cut surfaces (Fig. 1c.). The entire cecum, approximately one meter in length, had everted into the proximal colon (Fig. 1d.) and the associated caecal

submucosa was severely edematous. The feces were loose and watery.

Microscopically, the walls and adjacent connective tissue of multiple portal venules and peri-portal areas were infiltrated by inflammatory cells, predominantly macrophages, plasma cells, lymphocytes and a few eosinophils. Portal triads and perisinusoidal spaces were expanded by deposition of a collagenous stroma (as evident by Masson's trichrome stain) and bile ducts were often multiple and tortuous, indicative of severe chronic portal and perisinusoidal fibrosis and biliary hyperplasia. The wall of the cecum was diffusely edematous and the submucosa was often infiltrated by moderate numbers of eosinophils. There was marked hemorrhage and edema throughout the submucosa. There were some microscopic hemorrhages and ulcerations of the mucosa. The colonic and the cecal epithelia were diffusely attenuated and absent in some areas. The muscularis mucosa was diffusely very thick. Multiple areas of the colonic submucosa and muscularis mucosa were infiltrated by macrophages, epithelioid cells, giant cells, lymphocytes and plasma cells (Fig. 2a).

In both the liver and the colon, the centers of granulomas often contained non-operculated, thin, brown-walled eggs with a terminal spine and containing a ciliated miracidium, consistent with eggs of *Schistosoma* sp. (Parris, Michie et al. 2014) (Fig. 2b and 2d).



Fig. 1: (a) Emaciated elephant carcass (b) Ascites (arrow head), (c) White, multifocal to coalescing portal fibrosis (arrow) and (d) Intussusception: the caecum (arrow) is everted into the lumen of proximal colon




Fig. 2: (a) Granulomatous phlebitis with intralesional *Schistosoma* eggs (arrow) (HE) (Bar = 100µm), (b) High power view of a *Schistosoma* egg with ciliated miracidium (HE) (Bar = 20µm), (c) Portal fibrosis (arrow) and biliary hyperplasia (arrow head) (Masson's trichrome stain) (Bar = 200 µm) and (d) *Schistosoma* eggs in the sub-mucosa of the colon (thin arrows) (HE) (Bar = 100µm)

The pathological findings described here are consistent with those of human hepatic and intestinal schistosomiasis (Andrade, 1987). Ascites likely resulted from portal hypertension caused by hepatic portal and perisinusoidal fibrosis, perhaps enhanced by protein malnutrition associated with the prolonged clinical illness. The inflammatory lesions of the colon and in the cecum were implicated as the cause to the cecocolic intussusception, similar to previously described cecocolic intussusception in horses with parasite-induced intestinal inflammation (Mair *et al.*, 2000).

The Asian elephant in Sri Lanka is categorized as an endangered species in the International Union for Conservation of Nature (IUCN) red list (Choudhury *et al.*, 2008) and its population is rapidly declining due to human-elephant conflicts and habitat destruction in South East Asia. Information on infectious and non-infectious diseases of this species is needed to better understand health issues that may pose additional risks to the species.

The schistosome trematode, *Bivitellobilharzia nairi*, infects Asian elephants (Rajapakse *et al.*, 2003; Vimalraj *et al.*, 2012; Rajapakse *et al.*, 2013) whereas *B. loxodontae* infects African elephants (Brant *et al.*, 2013). This paper contributes to understanding health issues and extends the limited literature of *Schistoma* infection in Asian and African elephants. To our knowledge, this is the first report of cecocolic intussusception in an Asian elephant associated with hepato-intestinal schistosomiasis.

Acknowledgments

We acknowledge Dr. F.A. Leighton, Dr. Oswin Perera and Dr. N.K. Jayasekara for their kind assistance with manuscript preparation. We also thank the staff members of the Elephant Orphanage, Pinnawala and the Division of Veterinary Pathology, University of Peradeniya, Sri Lanka. This study received funding from Canada's International Development Research Centre (IDRC) through the project entitled "Building Research Excellence in Wildlife and Human Health in Sri Lanka", which is being implemented by the partner organizations of the Sri Lanka Wildlife

Health Centre and the Canadian Wildlife Health Cooperative.

References

- Agatsuma T, Rajapakse RPVJ, Kuruwita VY, Iwagami M and Rajapakse RC (2004) Molecular taxonomic position of the elephant schistosome, *Bivitellobilharzia nairi*, newly discovered in Sri Lanka. *Parasitology International* **53**(1): 69-75. DOI: 10.1016/j.parint.2003.11.003
- Andrade ZA (1987) Pathology of human schistosomiasis. *Memórias do Instituto Oswaldo Cruz* **82**: 17-23. DOI: 10.1590/S0074-02761987000800005
- Brant SV, Pomajbíková K, Modry D, Petrželková KJ, Todd A and Loker ES (2013) Molecular phylogenetics of the elephant schistosome *Bivitellobilharzia loxodontae* (Trematoda: Schistosomatidae) from the Central African Republic. *Journal of helminthology* **87**(01): 102-107. DOI: 10.1017/S0022149X1200003X
- Choudhury A, Choudhury DKL, et al. (2008) *Elephas maximus*. *The IUCN Red List of Threatened Species. Version 2014.3*.
- Mair TS, Sutton DGM and Love S (2000) Caecocolic and caecocolic intussusceptions associated with larval cyathostomosis in four young horses. *Equine Veterinary Journal* **32**(S32) : 77-80. DOI : 10.1111/j.2042-3306.2000.tb05339.x
- Parris V, Michie K, Andrews T, Nsutebu EF, Squire SB, Miller AR and Beadsworth MB (2014) Schistosomiasis japonicum diagnosed on liver biopsy in a patient with hepatitis B co-infection: a case report. *Journal of medical case reports* **8**(1): 45. DOI: 10.1186/1752-1947-8-45
- Rajapakse, R. P. V. J., Iwagami, M., Wickramasinghe, S., Walker, S. M., & Agatsuma, T. (2013). Morphology and surface topography of the schistosome *Bivitellobilharzia nairi* from the Asian elephant (*Elephas maximus maximus*) in Sri Lanka. *Journal of helminthology*, **87**(03), 348-355. DOI: 10.1017/S0022149X12000417
- Vimalraj PG, Jayathangaraj MG, Sridhar R, Senthilkumar TMA and Latchumikanthan A (2012) Elephant schistosome (*Bivitellobilharzia nairi*) in free-ranging Asian elephants (*Elephas maximus*) of Sathyamangalam forest division of Tamil Nadu State. *J. Vet. Parasitol.* **26**: 80-81.